

CP Violation in Two-Body Hadronic Decays of B_c Meson ^{*}

Youshan Dai ^{a,b} and Dongsheng Du ^a

a. Institute of High Energy Physics, Chinese Academy of Sciences

P.O.Box 918(4), Beijing, 100039, P.R.China

b. Department of Physics, Hangzhou University, Hangzhou, 310028, P.R.China

Using the next-to-leading order low energy effective Hamiltonian, the CP asymmetries for the B_c meson decays into meson pair are calculated in the spectator approximation. We do not compute the hadronic matrix elements directly, instead, we use the amplitude ratios to estimate the CP asymmetries. This is quite different from the previous works in the literature. The values of the momentum squared carried by the virtual particles in timelike penguin diagrams are also carefully discussed. From our calculated results, the best decay modes to observe CP violation in B_c decays would be $B_c^- \rightarrow \bar{D}^{*0} K^{*-}$, $\bar{D}^0 K^{*-}$, $\bar{D}^{*0} K^-$, $\bar{D}^0 K^-$ and $B_c^- \rightarrow \eta_c D^-$, which need about 10^8 of $B_c^\pm$ events in experiment.

PACS numbers: 11.30.Er, 13.20.He, 13.25.Hw, 13.25.-k

^{*}Supported in part by National Natural Science Foundation of China.

I. INTRODUCTION

One of the main aims of B factories is the observation of CP violation. The B_u, B_d and B_s meson decays and CP violation have been discussed extensively. The decays of B_c meson ($\bar{b}c$ and $b\bar{c}$ bound states) seems to be another valuable window for probing the origin of CP violation. Since large number of B_c mesons is expected to be produced at hadronic colliders like LHC or Tevatron [1], to examine the features of the B_c meson decays and CP violation becomes more and more interesting for both experimental efforts and theoretical studies.

Theoretical predictions about B_c meson decays are made in many previous works with different models [2]. The results are strongly model-dependent. It is difficult to judge these different results for lack of experimental data for the time being. There are also several works which pay attention to CP violation in B_c meson decays. In Ref.[3] the BSW mode and in Ref.[4] the Bethe-Salpeter formulism are used to calculate CP violating asymmetries for two-body mesonic decays of the B_c meson. There are many uncertainties due to the model-dependence and different choice of parameter values .

It is well known that there is only direct CP violation in B_c meson decays which requires that two decay amplitudes must have both different weak phases and different strong phases. The weak phases come from the CKM matrix. According to the new experimental results, the favorite values of the Wolfenstein parameters for the CKM matrix are changed, the main difference compared with the old ones is the sign of the parameter ρ [5]. Obviously, the new Wolfenstein parameter will influence the weak phases and the calculated results of the CP violating asymmetry. If we ignore soft strong phases which are difficult to estimate at present, hard strong phases can be estimated perturbatively using the low effective Hamiltonian and are relevant to the loop-integral functions of penguin diagrams. In general, decay amplitudes and the CP violating asymmetries rely on hadronic matrix elements which are strongly model-dependent, but in the spectator approximation (i.e. ignore the contributions of the annihilation and spacelike penguin diagrams, which are believed to be form factor and color suppressed), CP violating asymmetries can be obtained without calculation of the hadronic matrix elements for many decay processes of the B_c meson. In this paper, unlike the previous

works in the literature [3,4], we try to estimate the CP violating asymmetry of the B_c decays into meson pair without calculating the hadronic matrix elements directly. So that part of the uncertainties caused by the direct computation of the hadronic matrix elements are avoided. Of course we still need to use factorization approximation to factorized out some coefficients of the decay amplitudes. Thus the factorization approximation will definitely cause uncertainties. Another uncertainty source is the unknown momentum squared carried by the virtual particles in penguin diagrams. In the literature, one used to pick up a special value of k^2 from $[0, m_b^2]$ or $[\frac{1}{4}m_b^2, \frac{1}{2}m_b^2]$ for all timelike penguin diagrams. Obviously, it is a rather rough approximation. We shall discuss this problem in our paper and try to avoid the drawback of taking arbitrary values for k^2 based on the valence-quark assumption. In this article, we use the next-to-leading order low energy effective Hamiltonian [6] to calculate the CP violating asymmetry for the B_c decays into PP, PV and VV mesons in the spectator approximation (P- pseudoscalar meson, V- vector meson). In Section II, we first list the formula for the CP violating asymmetry. The quark-diagram amplitudes using the next-to-leading order low energy effective Hamiltonian are given in Section III. In Section IV, we discuss the value of k^2 carried by the virtual particles for the timelike penguin diagrams. The calculated results of the CP violating asymmetries for the B_c decays into meson pair are given in Section V. Finally Section VI is devoted to discussion and conclusion.

II. FORMULA OF CP ASYMMETRY

For the B_c meson decays to a final state f , we may, without loss of generality, write the decay transition amplitude as

$$A(f) = G_1 T_1 + G_2 T_2 \quad (2.1)$$

where G_1, G_2 are both multiplication of the CKM matrix elements. Assuming CPT invariance, the CP-conjugated amplitude is

$$\bar{A}(\bar{f}) = G_1^* T_1 + G_2^* T_2 \quad (2.2)$$

With the help of (2.1) and (2.2), the CP violating asymmetry can be written as

$$\mathcal{A}_{cp}(f) \equiv \frac{|A(f)|^2 - |\bar{A}(\bar{f})|^2}{|A(f)|^2 + |\bar{A}(\bar{f})|^2} = \frac{-2(ImG_1G_2^*)(ImT_1T_2^*)}{|G_1T_1|^2 + |G_2T_2|^2 + 2(ReG_1G_2^*)(ReT_1T_2^*)} \quad (2.3)$$

Define $G_1 = |G_1|e^{i\theta_1}$, $G_2 = |G_2|e^{i\theta_2}$ and $T_1 = |T_1|e^{i\delta_1}$, $T_2 = |T_2|e^{i\delta_2}$, where θ_1, θ_2 are the weak phases and δ_1, δ_2 are the strong phases respectively, then we have

$$\mathcal{A}_{cp}(f) = \frac{-2\sin(\theta_1 - \theta_2)\sin(\delta_1 - \delta_2)}{\left|\frac{G_1T_1}{G_2T_2}\right| + \left|\frac{G_2T_2}{G_1T_1}\right| + 2\cos(\theta_1 - \theta_2)\cos(\delta_1 - \delta_2)} \quad (2.4)$$

Obviously, the decay amplitude must have both different weak phases ($\theta_1 \neq \theta_2$) coming from the CKM matrix and different strong phases ($\delta_1 \neq \delta_2$) coming from final state interactions (FSI), otherwise there are no CP violation. Introduce another angle ζ which is defined as

$$h \equiv \left|\frac{G_1T_1}{G_2T_2}\right| \equiv tg\frac{\zeta}{2}, \quad \sin\zeta = \frac{2h}{1+h^2} > 0 \quad (2.5)$$

then the CP violating asymmetry formula becomes

$$\begin{aligned} \mathcal{A}_{cp}(f) &= \frac{\mathcal{A}_0}{1+\mathcal{A}_1} \\ \mathcal{A}_0 &= -\sin\zeta\sin(\theta_1 - \theta_2)\sin(\delta_1 - \delta_2) \\ \mathcal{A}_1 &= \sin\zeta\cos(\theta_1 - \theta_2)\cos(\delta_1 - \delta_2) \end{aligned} \quad (2.6)$$

We can see that the magnitude of the direct CP violation depends on three angles: weak phase ($\theta_1 - \theta_2$), strong phase ($\delta_1 - \delta_2$) and the angle ζ .

(i) The weak phase ($\theta_1 - \theta_2$) is decided by the CKM matrix elements, $\sin(\theta_1 - \theta_2) = \frac{ImG_1G_2^*}{|G_1G_2|}$, $\cos(\theta_1 - \theta_2) = \frac{ReG_1G_2^*}{|G_1G_2|}$. According to the Wolfenstein representation of the CKM matrix [7]

$$V = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} = \begin{pmatrix} 1 - \frac{\lambda^2}{2} & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1 - \frac{\lambda^2}{2} & A\lambda^2 \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 \end{pmatrix} + O(\lambda^4) \quad (2.7)$$

For two-body mesonic B_c^- decays, if we only consider b quark decay, we have, $|G_1G_2| = |V_{ud}^*V_{cd}^*V_{ub}V_{cb}| = |V_{us}^*V_{cs}^*V_{ub}V_{cb}| \approx A^2\lambda^6(1 - \frac{\lambda^2}{2})\sqrt{\rho^2 + \eta^2}$, and $ImG_1G_2^* = \pm J \approx \pm A^2\lambda^6(1 - \frac{\lambda^2}{2})\eta$. It follows that

$$\sin(\theta_1 - \theta_2) = \pm \frac{\eta}{\sqrt{\rho^2 + \eta^2}}, \quad |\theta_1 - \theta_2| = \gamma \quad (2.8)$$

where the angle γ is just one of the angles of the unitarity triangle for the CKM matrix elements in Fig.1. From (2.8), we see that the weak phase $(\theta_1 - \theta_2)$ is only decided by the Wolfenstein parameters (ρ, η) .

(ii) The strong phase $(\delta_1 - \delta_2)$ is caused by final state interactions, in which the penguin effects (hard strong phases) can be estimated perturbatively and the rescattering effects (soft strong phases) is unknown at present. In this paper we will ignore the rescattering effects and discuss the penguin effects only.

(iii) The angle ζ relies on the ratio $h = |\frac{G_1 T_1}{G_2 T_2}| = tg \frac{\zeta}{2}$. Although the decay amplitudes T_1 and T_2 both can be calculated, but there are uncertainties since the calculation of hadronic matrix elements, therefore T_1, T_2 , are model dependent. However, for many B_c decay processes, we do not need to compute the decay amplitudes (hadronic matrix elements) individually because the ratio $\frac{T_1}{T_2}$ is independent of the hadronic matrix elements (in factorization approximation). For example, in our formula of the CP violating asymmetry, it only needs to focus on calculating the ratio h and $\sin \zeta = f(h) = \frac{2h}{1+h^2}$. In the case of $h \ll 1$ (or $h \gg 1$), we have $\sin \zeta = f(h) \approx 2h$ (or $\frac{2}{h}$) $\ll 1$, $|\mathcal{A}_1| \ll 1$, so the CP violating asymmetry (2.6) can be simplified as

$$\mathcal{A}_{cp}(f) = \frac{\mathcal{A}_0}{1 + \mathcal{A}_1} \approx \mathcal{A}_0 = -\sin \zeta \sin(\theta_1 - \theta_2) \sin(\delta_1 - \delta_2) \quad (for \ h \ll 1 \ or \ h \gg 1) \quad (2.9)$$

III. THE QUARK-DIAGRAM AMPLITUDES

Following Ref.[6], the next-to-leading order low energy effective Hamiltonian for $|\Delta B|=1$ is given at the renormalization scale $\mu \sim m_b$ as

$$\mathcal{H}_{eff}(|\Delta B|=1) = \frac{G_F}{\sqrt{2}} \sum_{\substack{q=u,c \\ q'=d,s}} V_{qq'} V_{qb}^* \left\{ C_1(\mu) Q_1 + C_2(\mu) Q_2 + \sum_{k=3}^{10} C_k(\mu) Q_k \right\} + h.c. \quad (3.1)$$

where the Wilson coefficients $C_i(\mu)$ ($i = 1, 2, \dots, 10$) are calculated in the renormalization group improved perturbation theory and include the leading and next-to-leading order QCD corrections. Q_1 and Q_2 are the tree diagram operators; $Q_3, \dots, Q_6$ are the QCD penguin

diagram operators; whereas $Q_7, \dots, Q_{10}$ are the electroweak penguin diagram operators.

$$\begin{aligned}
Q_1 &= (\bar{q}'_\alpha q_\beta)_{V-A} (\bar{q}_\beta b_\alpha)_{V-A} & Q_2 &= (\bar{q}' q)_{V-A} (\bar{q} b)_{V-A} \\
Q_{3(5)} &= (\bar{q}' b)_{V-A} \sum_{q''} (\bar{q}'' q'')_{V-A(V+A)} & Q_{4(6)} &= (\bar{q}'_\alpha b_\beta)_{V-A} \sum_{q''} (\bar{q}''_\beta q''_\alpha)_{V-A(V+A)} \\
Q_{7(9)} &= \frac{3}{2} (\bar{q}' b)_{V-A} \sum_{q''} e_{q''} (\bar{q}'' q'')_{V+A(V-A)} & Q_{8(10)} &= \frac{3}{2} (\bar{q}'_\alpha b_\beta)_{V-A} \sum_{q''} e_{q''} (\bar{q}''_\beta q''_\alpha)_{V+A(V-A)}
\end{aligned} \tag{3.2}$$

In (3.2), q'' is running over the quark flavors being active at the scale $\mu \sim m_b$ ($q'' \in u, d, s, c, b$), $e_{q''}$ are the corresponding quark charges in unite of $|e|$, α and β are $SU(3)_c$ color indices, $(V \pm A)$ refer to $\gamma_\mu(1 \pm \gamma_5)$. In general, the nonleptonic B_c decays can occur through both the spectator channels and nonspectator channels. The latter are difficult to deal with at present and are commonly assumed to be form factor suppressed compared to the former ones. In spectator approximation, the two-body mesonic B_c^- decay can be described by quark diagrams as in Fig.2. Using $\mathcal{H}_{eff}$ (3.1) in a renormalization-scheme independent way [8] and factorization approximation, the quark-diagram amplitudes of Fig.2.(where the CKM matrix elements have been singled out) can be obtained

$$\begin{aligned}
T(a^{tree}) &= a_1 A & T(a_q^{pen}) &= [(a_3 + a_9) + \xi_f(a_5 + a_7) + (1 + \xi_f)b_1 G_q(k^2)] A \\
T(b^{tree}) &= a_2 B & T(b_q^{pen}) &= [(a_3 - \frac{1}{2}a_9) + \xi_f(a_5 - \frac{1}{2}a_7) + (1 + \xi_f)b_2 G_q(k^2)] B
\end{aligned} \tag{3.3}$$

where $T(a^{tree})$ refer to the amplitude of the diagram a^{tree} in Fig.2. etc., A and B denote the factorized hadronic matrix elements

$$\begin{aligned}
A &\equiv \frac{G}{\sqrt{2}} \langle X^- | (\bar{q}_1 q_2)_{V-A} | 0 \rangle \langle X^0 | (\bar{q}_3 b)_{V-A} | B_c^- \rangle & for X^- = (q_1 \bar{q}_2) \quad , \quad X^0 = (q_3 \bar{c}) \\
B &\equiv \frac{G}{\sqrt{2}} \langle X^0 | (\bar{q}_1 q_2)_{V-A} | 0 \rangle \langle X^- | (\bar{q}_3 b)_{V-A} | B_c^- \rangle & for X^0 = (q_1 \bar{q}_2) \quad , \quad X^- = (q_3 \bar{c})
\end{aligned} \tag{3.4}$$

In (3.3), ξ_f arises from the transformation of (V-A)(V+A) currents into (S+P)(S-P) and further into (V-A)(V-A) ones for Q_{5-8}

$$\xi_f = \begin{cases} \frac{(1+\eta_{X^-})M_{X^-}^2}{(m_{q'}+m_{q_v})(\eta_{X^0}m_b-m_{q_v})} & for a^{pen} \text{ diagrams} \\ \frac{(1+\eta_{X^0})M_{X^0}^2}{(m_{q'}+m_{q'_v})(\eta_{X^-}m_b-m_{q'_v})} & for b^{pen} \text{ diagrams} \end{cases} \tag{3.5}$$

where

$$\eta_{X^0}(or \eta_{X^-}) = \begin{cases} +1 & for X^0(or X^-) = P \text{ (pseudoscalar meson)} \\ -1 & for X^0(or X^-) = V \text{ (vector meson)} \end{cases} \tag{3.6}$$

From (3.5) and (3.6), we can see that $\xi_f = 0$ when X^- is vector meson in a^{pen} diagrams and when X^0 is vector meson in b^{pen} diagrams. The reason is the following: using the Fierz rearrangements for Q_{5-8} , the calculated results of the decay amplitudes are proportional to the factor $P^\mu(X^-)\epsilon_\mu(X^-)$ (or $P^\mu(X^0)\epsilon_\mu(X^0)$) which is equal to zero (P^μ and ϵ_μ are momentum and polarization vectors of the vector meson respectively). In (3.3), a_k is defined as

$$a_{2i-1} \equiv \frac{\bar{c}_{2i-1}}{N_c} + \bar{c}_{2i} \quad a_{2i} \equiv \frac{\bar{c}_{2i}}{N_c} + \bar{c}_{2i-1} \quad (i = 1, 2, 3, 4, 5) \quad (3.7)$$

where $\bar{c}_k$ ($k = 1, \dots, 10$) are the renormalization scheme independent Wilson coefficients $\bar{c}_k(\mu)$ at $\mu \sim m_b$. We use $\alpha_s(m_Z) = 0.118$, $\alpha_{em}(m_Z) = \frac{1}{128}$, $m_t = 174 GeV$, $\bar{c}_i$ is taken as [9]

$$\begin{aligned} \bar{c}_1 &= -0.313 & \bar{c}_2 &= 1.150 \\ \bar{c}_3 &= 0.017 & \bar{c}_4 &= -0.037 & \bar{c}_5 &= 0.010 & \bar{c}_6 &= -0.047 \\ \bar{c}_7 &= -0.001\alpha_{em} & \bar{c}_8 &= 0.049\alpha_{em} & \bar{c}_9 &= -1.321\alpha_{em} & \bar{c}_{10} &= 0.267\alpha_{em} \end{aligned} \quad (3.8)$$

and

$$b_1 = \frac{2}{3} \left[\frac{\alpha_s}{8\pi} \bar{c}_2 \left(1 - \frac{1}{3N_c}\right) + \frac{\alpha_{em}}{3\pi} \left(\bar{c}_1 + \frac{\bar{c}_2}{3}\right) \frac{1}{N_c} \right] \quad (3.9)$$

$$b_2 = \frac{2}{3} \left[\frac{\alpha_s}{8\pi} \bar{c}_2 \left(1 - \frac{1}{3N_c}\right) - \frac{\alpha_{em}}{6\pi} \left(\bar{c}_1 + \frac{\bar{c}_2}{3}\right) \frac{1}{N_c} \right]$$

$$G_q(k^2) = \frac{3}{2} \left[\frac{10}{9} - F_q(k^2) \right] \quad (3.10)$$

where $F_q(k^2)$ denotes the penguin loop-integral function with momentum transfer squared k^2 at the scale $\mu \sim m_b$

$$F_q(k^2) = -4 \int_0^1 dx x(1-x) \ln \left[\frac{m_q^2 - x(1-x)k^2}{m_b^2} \right] \quad (3.11)$$

Supposing $k^2 > 4m_q^2$ ($q = u, c$), the parameter $r_q = \frac{4m_q^2}{k^2} < 1$, the function $G_q(k^2)$ can be analytically expressed as [10]

$$G_q(k^2) = \ln \frac{m_q^2}{m_b^2} - r_q + \left(1 + \frac{r_q}{2}\right) \sqrt{1 - r_q} \ln \frac{1 + \sqrt{1 - r_q}}{1 - \sqrt{1 - r_q}} + i\pi \left(1 + \frac{r_q}{2}\right) \sqrt{1 - r_q} \quad (3.12)$$

In the case of $q = u$, since $r_u \ll 1$, we have $G_u(k^2) \approx \ln \frac{k^2}{m_b^2} + i\pi$, the absorptive part of the $G_q(k^2)$ will lead to direct CP violation.

IV. MOMENTUM SQUARED CARRIED BY THE VIRTUAL PARTICLES

The penguin loop-integral function $F_q(k^2)$ depends crucially on the k^2 carried by the virtual gluon, photon and Z^0 . In the literature, one used to pick up a special fixed value of k^2 for timelike penguin diagrams from $[0, m_b^2]$ or $[\frac{1}{4}m_b^2, \frac{1}{2}m_b^2]$ [10], but in general, taking the same value of k^2 for all different quark diagrams is not correct. This problem has been discussed in Ref.[11], we shall further discuss it carefully in the following. For timelike penguin diagram of the two-body mesonic decay $B_c^- \rightarrow XY$ as illustrated in Fig.3, using the 4-momentum conservation ($\bar{c}$ as spectator quark), we have

$$\begin{cases} k^2 = m_b^2 + m_q^2 - 2E_b E_q + 2 \vec{p}_b \cdot \vec{p}_q \\ \vec{p}_b = \vec{p}_q + \vec{p}_{\bar{q}_v} + \vec{p}_{q_v} \\ E_b = E_q + E_{\bar{q}_v} + E_{q_v} \end{cases} \quad (4.1)$$

In the rest frame of the X meson, $\vec{p}_X = \vec{p}_q + \vec{p}_{\bar{q}_v} = 0$. Denote $a \equiv E_q + E_{\bar{q}_v}$, from (4.1), we have

$$\begin{cases} |\vec{p}_b| = |\vec{p}_{q_v}| \\ E_b = a + E_{q_v} \end{cases} \quad \begin{cases} |\vec{p}_q| = |\vec{p}_{\bar{q}_v}| \\ E_q = a - E_{\bar{q}_v} \end{cases} \quad (4.2)$$

then we obtain

$$\begin{aligned} E_b &= \frac{1}{2a}(m_b^2 + a^2 - m_{q_v}^2) \quad , \quad |\vec{p}_b| = \frac{1}{2a}\sqrt{[(m_b + a)^2 - m_{q_v}^2][(m_b - a)^2 - m_{q_v}^2]} \\ E_q &= \frac{1}{2a}(m_q^2 + a^2 - m_{q_v}^2) \quad , \quad |\vec{p}_q| = \frac{1}{2a}\sqrt{[(m_q + a)^2 - m_{q_v}^2][(m_q - a)^2 - m_{q_v}^2]} \end{aligned} \quad (4.3)$$

Because $q, \bar{q}_v$ form a bound state X, in the valence-quark assumption, there are $p_q = \eta_q p_X + p$, $p_{\bar{q}_v} = \eta_{\bar{q}_v} p_X - p$, where $\eta_q = \frac{m_q}{m_q + m_{q_v}}$, $\eta_{\bar{q}_v} = \frac{m_{q_v}}{m_q + m_{q_v}}$; p_X is the total 4-momentum of the bound state X and p is the relative 4-momentum of q vs. $\bar{q}_v$ inside the X meson. In the rest frame of the X meson, it is easy to get $E_q = \frac{1}{2M_X}(m_q^2 + M_X^2 - m_{q_v}^2)$. So we have $a = M_X$. In eq.(4.1), taking $\vec{p}_b \cdot \vec{p}_q = |\vec{p}_b| |\vec{p}_q| \cos\varphi$, where φ is the angle between $\vec{p}_b$ and $\vec{p}_q$ in the rest frame of the X meson, $\varphi = \angle(\vec{p}_b, \vec{p}_q)$, then we get

$$k^2(\varphi) = m_b^2 + m_q^2 - 2E_b E_q + 2|\vec{p}_b| |\vec{p}_q| \cos\varphi \quad , \quad \bar{k}^2 = \frac{1}{\pi} \int_0^\pi d\varphi k^2(\varphi) = m_b^2 + m_q^2 - 2E_b E_q \quad (4.4)$$

$$(k^2)_{max} = m_b^2 + m_q^2 - 2E_b E_q + 2|\vec{p}_b| |\vec{p}_q| \quad , \quad (k^2)_{min} = m_b^2 + m_q^2 - 2E_b E_q - |\vec{p}_b| |\vec{p}_q| \quad (4.5)$$

where E_b , E_q , $|\vec{p}_b|$, $|\vec{p}_q|$ can be obtained from eq.(4.3) by taking $a = M_X$. Accordingly we find that the average value of the k^2 is

$$\frac{\bar{k}^2}{m_b^2} = \frac{1}{2} \left[1 + \frac{m_{q_v}^2 - m_q^2}{M_X^2} \left(1 - \frac{m_{q_v}^2}{m_b^2} \right) + \frac{m_q^2 + 2m_{q_v}^2 - M_X^2}{m_b^2} \right] \quad (4.6)$$

For the timelike penguin diagram, $0 \leq (k^2)_{min} \leq k^2(\varphi) \leq (k^2)_{max} \leq m_b^2$. Since φ is unknown in experiment, we take the average value of the $\bar{k}^2$ to calculate the penguin loop-integral function. Taking the quark masses $(m_u, m_d, m_s, m_c, m_b) = (0.005, 0.01, 0.175, 1.35, 4.8)$ GeV and using the meson masses M_X according to the particle data [12], the values $\frac{\bar{k}^2}{m_b^2}$ of the timelike penguin diagram for different B_c^- decay processes are given in Table II.

V. CP ASYMMETRY

As discussed in Section II, the CP violation asymmetry of B_c meson decays into meson pair depends on the three angles: weak phase $(\theta_1 - \theta_2)$, strong phase $(\delta_1 - \delta_2)$ and the angle ζ . The weak phase $(\theta_1 - \theta_2)$ is determined by the Wlffenstein parameter (ρ, η) , the strong phase $(\delta_1 - \delta_2)$ and the angle ζ are decided by the quark-diagram amplitudes. For many decay processes, there is no need to calculate the hadronic matrix elements because in the factorization approximation, the factorized quark-diagram amplitudes depend only on a single (dominant) hadronic matrix element (A or B) which is cancelled in the ratio $\frac{T_1}{T_2}$.

In below we discuss the decay process $B_c^- \rightarrow \bar{D}^0 \pi^-$ as an example to illustrate how to calculate the CP violating asymmetry in our method. In the spectator approximation, the decay amplitude for the $B_c^- \rightarrow \bar{D}^0 \pi^-$ is

$$A(f) = G_1 T_1 + G_2 T_2 = V_{ud}^* V_{ub} [T(a^{tree}) + T(a_u^{pen})] + V_{cd}^* V_{cb} T(a_c^{pen}) \quad (5.1)$$

using the quark-diagram amplitude calculated in section.III, we get

$$\frac{T_1}{T_2} = \left| \frac{T_1}{T_2} \right| e^{i(\delta_1 - \delta_2)} = \frac{a_1 + a_3 + a_9 + \xi_{(\bar{D}^0 \pi^-)}(a_5 + a_7) + [1 + \xi_{(\bar{D}^0 \pi^-)}] b_1 G_u(\bar{k}^2)}{a_3 + a_9 + \xi_{(\bar{D}^0 \pi^-)}(a_5 + a_7) + [1 + \xi_{(\bar{D}^0 \pi^-)}] b_1 G_c(k^2)} \quad (5.2)$$

$$\begin{cases} \sin(\delta_1 - \delta_2) = \frac{1}{|T_1 T_2|} (Im T_1 Re T_2 - Im T_2 Re T_1) \\ \cos(\delta_1 - \delta_2) = \frac{1}{|T_1 T_2|} (Re T_1 Re T_2 + Im T_1 Im T_2) \end{cases} \quad (5.3)$$

where $\xi_{(\bar{D}^0\pi^-)} = \frac{2m_{\pi^-}^2}{(m_d+m_u)(m_b-m_u)} = 0.545$, for $B_c^- \rightarrow \bar{D}^0\pi^-$, $\frac{k^2}{m_b^2} = 0.498$, $G_u(\bar{k}^2) = -0.697 + i\pi$, $G_c(\bar{k}^2) = -2.059 + i2.500$. Taking $N_c = N_c^{eff} = 2$, which is favored by experimental data [13], then the numerical results can be obtained as $\left|\frac{T_1}{T_2}\right| = 14.55$, $\sin(\delta_1 - \delta_2) = -0.197$, $\cos(\delta_1 - \delta_2) = -0.980$. From the Wolfenstein representation of the CKM matrix, $G_1 = V_{ud}^*V_{ub} = A\lambda^3(1 - \frac{\lambda^2}{2})(\rho - i\eta)$, $G_2 = V_{cd}^*V_{cb} = -A\lambda^3$, it follows that

$$\begin{cases} \sin(\theta_1 - \theta_2) = \frac{ImG_1G_2^*}{|G_1G_2|} = \frac{\eta}{\sqrt{\rho^2 + \eta^2}} \\ \cos(\theta_1 - \theta_2) = \frac{ReG_1G_2^*}{|G_1G_2|} = \frac{-\rho}{\sqrt{\rho^2 + \eta^2}} \end{cases} \quad \begin{cases} h = \left|\frac{G_1T_1}{G_2T_2}\right| = \frac{\lambda^2\sqrt{\rho^2 + \eta^2}}{(1 - \frac{\lambda^2}{2})} \left|\frac{T_1}{T_2}\right| \\ \sin\zeta = \frac{2h}{1+h^2} \end{cases} \quad (5.4)$$

According to our CP asymmetry formula derived in section.II, we have $\mathcal{A}_{cp} = \frac{\mathcal{A}_0}{1+\mathcal{A}_1}$, where $\mathcal{A}_0 = -\sin\zeta\sin(\theta_1 - \theta_2)\sin(\delta_1 - \delta_2)$, $\mathcal{A}_1 = \sin\zeta\cos(\theta_1 - \theta_2)\cos(\delta_1 - \delta_2)$. In this paper, we take $\lambda = 0.220$, $\eta = 0.336$, $\rho = 0.160$ (the value of the Wolfenstein parameter A has no effect on our calculation) [5]. For numerical estimation, the results are $\sin(\theta_1 - \theta_2) = 0.903$, $\cos(\theta_1 - \theta_2) = -0.430$, $\sin\zeta = 0.365$ and $\mathcal{A}_0 = 0.0649$, $\mathcal{A}_1 = 0.154$, $\mathcal{A}_{cp}(f = \bar{D}^0\pi^-) = 5.62 \times 10^{-2}$.

The quark-diagram amplitudes for the B_c^- decays into PP, PV, and VV mesons in the spectator approximation are given in Table.I. It is seen that we only need to compute the ratio $\frac{T_1}{T_2}$ and there is no need to calculate the hadronic matrix elements directly in many cases, but for the B_c^- meson decays to $(c\bar{c})D^-$, $(c\bar{c})D^{*-}$, $(c\bar{c})D_s^-$, $(c\bar{c})D_s^{*-}$, the hadronic matrix elements A and B still appear in the ratio

$$\frac{T_1}{T_2} = \frac{a_3 + a_9 + \xi_f(a_5 + a_7) + (1 + \xi_f)b_1G_u(\bar{k}^2)}{a_1 + a_2\frac{B}{A} + a_3 + a_9 + \xi_f(a_5 + a_7) + (1 + \xi_f)b_1G_c(\bar{k}^2)} \quad (5.5)$$

Fortunately a_2 is colour suppressed compared with a_1 and $\frac{B}{A} \sim O(1)$, so the calculated results are not sensitive to the value of $\frac{B}{A}$. In this paper, we simply take $\frac{B}{A} = 1$ for the estimation of CP asymmetries of these B_c^- decay processes.

All calculated results of the CP violating asymmetries for two-body mesonic decays of the B_c^- meson are presented in Table II.

VI. DISCUSSION AND CONCLUSION

Using the next-to-leading order low energy effective Hamiltonian and the quark-diagram amplitude method, we have calculated the CP violating asymmetries for two-body mesonic

decays of the B_c meson. For most decay processes (except for $B_c^- \rightarrow (c\bar{c})D^-$, $(c\bar{c})D^{*-}$, $(c\bar{c})D_s^-$, $(c\bar{c})D_s^{*-}$), the CP violating asymmetries do not rely on the hadronic matrix elements since they are canceled in the ratio $\frac{T_1}{T_2}$. So there is no model-dependence caused by calculating the hadronic matrix elements.

The CP violating asymmetry is proportional to $\mathcal{A}_0 = -\sin\zeta \sin(\theta_1 - \theta_2) \sin(\delta_1 - \delta_2)$. From Table II, we see that $\sin(\theta_1 - \theta_2)$ is the same for all decay processes (except for a sign) which is determined by the Wolfenstein parameters (ρ , η) only. In the processes (1)-(16), $\sin(\theta_1 - \theta_2) = \frac{\eta}{\sqrt{\rho^2 + \eta^2}} = 0.903$, $\cos(\theta_1 - \theta_2) = \frac{-\rho}{\sqrt{\rho^2 + \eta^2}} = -0.430$, $\left| \frac{G_1}{G_2} \right| = (1 - \frac{\lambda^2}{2})\sqrt{\rho^2 + \eta^2}$; while in processes (17)-(30), $\sin(\theta_1 - \theta_2) = \frac{-\eta}{\sqrt{\rho^2 + \eta^2}}$, $\cos(\theta_1 - \theta_2) = \frac{\rho}{\sqrt{\rho^2 + \eta^2}}$, $\left| \frac{G_1}{G_2} \right| = \frac{\lambda^2 \sqrt{\rho^2 + \eta^2}}{(1 - \frac{\lambda^2}{2})}$ which is doubly Cabibbo suppressed. In processes (1)-(6) and (17)-(20), T_1 is dominated by the tree diagram and T_2 only by the penguin diagram only, so $|T_1| \gg |T_2|$; in processes (9)-(12) and (21)-(26), T_1 and T_2 are both from the penguin diagram contribution only, so $|T_1| \sim |T_2|$; while in the processes (13)-(16) and (27)-(30), T_2 is dominated by the tree diagram and T_1 by the penguin diagram, so $|T_2| \gg |T_1|$. Since the CP violating asymmetry is proportional to $\sin\zeta = \frac{2h}{1+h^2}$, where $h = \left| \frac{G_1 T_1}{G_2 T_2} \right|$, if $h \gg 1$ or $h \ll 1$, $\sin\zeta$ is small. From Table II, The CP violating asymmetry is suppressed strongly by the small $\sin\zeta$ in the processes $B_c^- \rightarrow \psi D^-$, $\eta_c D_s^-$, ψD_s^- , $\eta_c D_s^{*-}$, ψD_s^{*-} . For the processes with $h \sim O(1)$ ($|G_1 T_1| \sim |G_2 T_2|$), $\sin\zeta$ is large, such as in the processes $B_c^- \rightarrow \pi^0 D^-$, $\rho^0 D^-$, $\rho^0 D^{*-}$, $\bar{D}^{*0} K^-$, $\bar{D}^0 K^{*-}$, $\bar{D}^{*0} K^{*-}$. The hard strong phase $(\delta_1 - \delta_2)$ is caused by the interference between penguin and tree diagrams or penguin diagram themselves, which rely on the penguin loop-integral functions $G_u(k^2)$ and $G_c(k^2)$. As discussed in Section IV, we take the average value of the momentum squared carried by virtual particles for our calculation. In most processes (below the $c\bar{c}$ threshold) $\frac{\bar{k}^2}{m_b^2} \sim 0.5$, but for producing $c\bar{c}$ pair ($B_c^- \rightarrow (c\bar{c})D^-$, $(c\bar{c})D^{*-}$, $(c\bar{c})D_s^-$, $(c\bar{c})D_s^{*-}$), $\frac{\bar{k}^2}{m_b^2} \sim 0.7$. From Table II, we can see that for the processes with interference of penguin diagram themselves only, such as the processes (9)-(12) and (21)-(26), $|\sin(\delta_1 - \delta_2)|$ is smaller than that for the processes with interference between penguin and tree diagrams. So, if the decay processes both satisfy: (i) $|G_1 T_1| \sim |G_2 T_2|$, (ii) with interference between penguin and tree diagrams, then the CP violating asymmetry will be large, such as the processes (5)-(8) and (17)-(20).

According to our CP violating asymmetry formula $\mathcal{A}_{cp} = \frac{\mathcal{A}_0}{1+\mathcal{A}_1}$, where $\mathcal{A}_0 = -\sin\zeta\sin(\theta_1 - \theta_2)\sin(\delta_1 - \delta_2)$, $\mathcal{A}_1 = \sin\zeta\cos(\theta_1 - \theta_2)\cos(\delta_1 - \delta_2)$, it can be seen that $\mathcal{A}_{cp}$ is proportional to $\mathcal{A}_0$ and the proportion factor is $\frac{1}{1+\mathcal{A}_1}$. If $|\mathcal{A}_1| \ll 1$, then $\mathcal{A}_{cp} \sim \mathcal{A}_0$. Since $\mathcal{A}_0 \propto \eta$, $\mathcal{A}_1 \propto \rho$, the CP violating asymmetry is more sensitive to the parameter η than the parameter ρ . In this paper, we have taken $\rho = 0.160$ [5] for numerical estimate, the main difference compare with the early literature is to change the sign of the parameter ρ . For the case of $|\mathcal{A}_1| \ll 1$, the influence of this change is very small, but in the case of $|\mathcal{A}_1| \sim 1$, it is important. In the processes (5)-(8), the CP violating asymmetry will be smaller than that with minus ρ ; while in the processes (17)-(20), it will be larger than that with minus ρ .

In this paper, we do not discuss the decay width of the B_c meson, since it needs to calculate the hadronic matrix elements which is strongly model-dependent. But for testing the CP violation in experiment, we need to find processes with both larger CP violating asymmetry and larger branching ratio. For three standard deviation (3σ) signature $\varepsilon_f N \sim \frac{9}{B_r \mathcal{A}_{cp}^2}$, where ε_f is the detecting efficiency of the final state, N is number of the $B_c^\pm$. Since lack of experimental data for the branching fraction of B_c decays at present, so in Table.II, we use the results of decay widths for the hadronic decay of the B_c^- calculated in Ref.[4] but take $\tau_{B_c}=0.5$ ps [14] to estimate the branching ratio B_r and $\varepsilon_f N$. This estimation show that in experiment the best decay modes to observe CP violation in B_c decays would be $B_c^- \rightarrow \bar{D}^{*0}K^{*-}$, $\bar{D}^0 K^{*-}$, $\bar{D}^{*0}K^-$, $\bar{D}^0 K^-$ and $B_c^- \rightarrow \eta_c D^-$, which need about 10^8 of $B_c^\pm$ events.

In summary, we have calculated the CP violating asymmetries for the B_c meson decays into PP, PV, VV mesons. It is only relevant to the three angles, weak phase $(\theta_1 - \theta_2)$, strong phase $(\delta_1 - \delta_2)$ and the angle ζ . If $|\sin(\delta_1 - \delta_2)|$ and $\sin\zeta$ are both large ($|\sin(\theta_1 - \theta_2)|$ is the same for all decay processes), the CP violating asymmetry would be large. There are less uncertainties in our calculated CP violating asymmetries since we avoided calculating the hadronic matrix elements in our method. In general, we need about 10^8 of $B_c^\pm$ events for testing the CP violation in experiment.

Finally we should mention that, in this paper, we only calculated the hard strong phases, the effects of the soft strong phases would also be important, it needs further study.

ACKNOWLEDGEMENTS

This work was supported in part by National Natural Science Foundation of China and Beijing Electron Positron Collider National Laboratory.

- [1] J.F.Donoghue, C.Ramirez, and G.Valencia, Phys.Rev.D **39**, 1947 (1989);
J.F.Donoghue and B.R.Holstein, Phys.Rev.D **40**, 238 (1989);
K.Cheung, Phys.Rev.Lett. **71**, 3413 (1993)
- [2] J.Gasser and H.Leutwyler, Phys.Rep. **87**, 77 (1982);
R.D.Peccei and J.Sola, Nucl.Phys.B **281**, 1 (1987);
D.S.Du and Z.Wang, Phys.Rev.D **39**, 1342 (1989);
J.Bijnens and E.de Rafael, Phys.Lett.B **273**, 483 (1991);
J.F.Donoghue, B.R.Holstein and D.Wyler, Phys.Rev.Lett, 3444 (1992); Phys.Rev.D **47**, 2089 (1993);
G.Ecker, J.Gasser, Nucl.Phys.B **490**, 239 (1997)
- [3] M.Masetti, Phys.Lett.B **286**, 160 (1992)
- [4] J.F.Liu and K.T.Chao, Phys.Rev.D **56**, 4133 (1997)
- [5] A.Ali, DESY 97-256, hep-ph/9801270;
F.Parodi, P. Roudeau and A.slocchi, hep-ph/9802289
- [6] A.J.Buras, M.Jamin, M.E.Lautenbacher and P.H.Weisz, Nucl.Phys.B **400**, 37 (1993)
A.J.Buras, M.Jamin and M.E.Lautenbacher, Nucl.Phys.B **444**, 75 (1993); ibid. B. **408**, 209 (1993)
- [7] L.Wolfenstein, Phys.Rev.Lett. **51**, 1945 (1983)
- [8] A.Buras, M.Jamin, M.Lautenbacher and P.Weisz, Nucl.Phys.B **370**, 69 (1992); ibid. **375**, 501 (1992)

- [9] N.G.Deshpande and X.G.He, Phys.Lett.B **336**, 471 (1994); Phys. Rev.Lett. **26**, 74 (1995)
- [10] J.M.Gérard and W.S.Hou, Phys.Rev.Lett. **62**, 855 (1989); Phys.Rev.D **43**, 2909 (1991)
- [11] D.S.Du and Z.Z.Xing, Phys.Lett.B **349**, 215 (1995)
- [12] Particle Data Group, Phys.Rev.D **54**, 454 (1996)
- [13] H.Y.Cheng and B.Tseng, Phys.Rev.D **51**, 6259 (1995)
- [14] M.Beneke, Fermilab-Pub-95/401-T

TABLE CAPTIONS

Table.I. The quark-diagram amplitudes for the B_c^- decays into PP, PV, VV mesons in the spectator approximation.

Table.II. The CP violating asymmetries for two-body mesonic decays of the B_c^- meson. where $h = \left| \frac{G_1 T_1}{G_2 T_2} \right|$, $\sin\zeta = \frac{2h}{1+h^2}$; $\mathcal{A}_{cp} = \frac{\mathcal{A}_0}{1+\mathcal{A}_1}$, $\mathcal{A}_0 = -\sin\zeta \sin(\theta_1 - \theta_2) \sin(\delta_1 - \delta_2)$, $\mathcal{A}_1 = \sin\zeta \cos(\theta_1 - \theta_2) \cos(\delta_1 - \delta_2)$. In processes (1)-(16), $\sin(\theta_1 - \theta_2)=0.903$, $\cos(\theta_1 - \theta_2)=-0.430$; while in processes (17)-(30), $\sin(\theta_1 - \theta_2)=-0.903$, $\cos(\theta_1 - \theta_2)=0.430$.

Table.I

No.	Final state f	The quark-diagram amplitudes
(1)	$\bar{D}^0 \pi^-$	$V_{ud}^* V_{ub} [T(a^{tree}) + T(a_u^{pen})] + V_{cd}^* V_{cb} T(a_c^{pen})$
(2)	$\bar{D}^{*0} \pi^-$	
(3)	$\bar{D}^0 \rho^-$	
(4)	$\bar{D}^{*0} \rho^-$	
(5)	$\pi^0 D^-$	$V_{ud}^* V_{ub} [T(b^{tree}) + T(b_u^{pen})] + V_{cd}^* V_{cb} T(b_c^{pen})$
(6)	$\pi^0 D^{*-}$	
(7)	$\rho^0 D^-$	
(8)	$\rho^0 D^{*-}$	
(9)	$K^0 D_s^-$	$V_{ud}^* V_{ub} T(b_u^{pen}) + V_{cd}^* V_{cb} T(b_c^{pen})$
(10)	$K^{*0} D_s^-$	
(11)	$K^{*0} D_s^{*-}$	
(12)	$K^{*0} D_s^{*-}$	
(13)	$\eta_c D^-$	$V_{ud}^* V_{ub} T(a_u^{pen}) + V_{cd}^* V_{cb} [T(a^{tree}) + T(b^{tree}) + T(a_c^{pen})]$
(14)	ψD^-	
(15)	$\eta_c D^{*-}$	
(16)	ψD^{*-}	
(17)	$\bar{D}^0 K^-$	$V_{us}^* V_{ub} [T(a^{tree}) + T(a_u^{pen})] + V_{cs}^* V_{cb} T(a_c^{pen})$
(18)	$\bar{D}^{*0} K^-$	
(19)	$\bar{D}^0 K^{*-}$	
(20)	$\bar{D}^{*0} K^{*-}$	
(21)	$\bar{K}^0 D^-$	$V_{us}^* V_{ub} T(b_u^{pen}) + V_{cs}^* V_{cb} T(b_c^{pen})$
(22)	$\bar{K}^{*0} D^-$	
(23)	$\bar{K}^{*0} D^{*-}$	
(24)	$\bar{K}^{*0} D^{*-}$	
(25)	ϕD_s^-	$V_{us}^* V_{ub} T(a_u^{pen}) + V_{cs}^* V_{cb} [T(a^{tree}) + T(b^{tree}) + T(a_c^{pen})]$
(26)	ϕD_s^{*-}	
(27)	$\eta_c D_S^-$	
(28)	ψD_s^-	
(29)	$\eta_c D_s^{*-}$	$V_{us}^* V_{ub} T(a_u^{pen}) + V_{cs}^* V_{cb} [T(a^{tree}) + T(b^{tree}) + T(a_c^{pen})]$
(30)	ψD_s^{*-}	

Table.II

No.	f	ξ_f	$\frac{k^2}{m_h^2}$	$\sin(\delta_1 - \delta_2)$	h	$\sin\zeta$	$\mathcal{A}_0$	$\mathcal{A}_1$	$\mathcal{A}_{cp} = \frac{\mathcal{A}_0}{1+\mathcal{A}_1}$	B_r	$\varepsilon_f N$
(1)	$\bar{D}^0\pi^-$	0.545	0.498	-0.197	5.284	0.365	0.0649	0.154	5.62×10^{-2}	2.68×10^{-6}	1.06×10^9
(2)	$\bar{D}^{*0}\pi^-$	-0.544	0.498	-0.274	27.93	0.0715	0.0177	0.0296	1.72×10^{-2}	2.18×10^{-6}	1.40×10^{10}
(3)	$\bar{D}^0\rho^-$	0	0.487	-0.202	9.010	0.219	0.0399	0.0922	3.65×10^{-2}	6.85×10^{-6}	9.86×10^8
(4)	$\bar{D}^{*0}\rho^-$	0	0.487	-0.202	9.010	0.219	0.0399	0.0922	3.65×10^{-2}	6.81×10^{-6}	9.92×10^8
(5)	$\pi^0 D^-$	0.380	0.500	-0.257	1.462	0.932	0.216	0.387	1.56×10^{-1}	2.28×10^{-7}	1.62×10^9
(6)	$\pi^0 D^{*-}$	-0.379	0.500	-0.316	5.528	0.350	0.0999	0.143	8.74×10^{-2}	7.17×10^{-8}	1.64×10^{10}
(7)	$\rho^0 D^-$	0	0.487	-0.255	2.401	0.710	0.163	0.295	1.26×10^{-1}	4.21×10^{-7}	1.35×10^9
(8)	$\rho^0 D^{*-}$	0	0.487	-0.255	2.401	0.710	0.163	0.295	1.26×10^{-1}	1.39×10^{-6}	4.08×10^8
(9)	$K^0 D_s^-$	0.577	0.557	-0.0616	0.3227	0.585	0.0325	-0.251	4.34×10^{-2}	2.38×10^{-7}	2.01×10^{10}
(10)	$K^0 D_s^{*-}$	-0.537	0.557	-0.189	0.3066	0.561	0.0957	-0.237	1.25×10^{-1}	2.67×10^{-8}	2.16×10^{10}
(11)	$K^{*0} D_s^-$	0	0.503	-0.0892	0.3161	0.575	0.0463	-0.246	6.14×10^{-2}	1.43×10^{-7}	1.67×10^{10}
(12)	$K^{*0} D_s^{*-}$	0	0.503	-0.0892	0.3161	0.575	0.0463	-0.246	6.14×10^{-2}	2.85×10^{-7}	8.38×10^9
(13)	$\eta_c D^-$	1.489	0.744	0.261	0.03055	0.0610	-0.0144	0.0253	1.40×10^{-2}	2.95×10^{-4}	1.56×10^8
(14)	ψD^-	-0.835	0.744	0.579	0.0007797	0.00156	-0.000816	-0.000547	8.16×10^{-4}	2.19×10^{-5}	6.17×10^{11}
(15)	$\eta_c D^{*-}$	0	0.699	0.286	0.01012	0.0202	-0.00522	0.00832	5.18×10^{-3}	6.89×10^{-5}	4.87×10^9
(16)	ψD^{*-}	0	0.699	0.286	0.01012	0.0202	-0.00522	0.00832	5.18×10^{-3}	4.95×10^{-4}	6.78×10^8
(17)	$\bar{D}^0 K^-$	0.565	0.433	-0.173	0.2579	0.484	-0.0756	-0.205	-9.51×10^{-2}	4.67×10^{-6}	2.13×10^8
(18)	$\bar{D}^{*0} K^-$	-0.564	0.433	-0.243	1.482	0.927	-0.203	-0.387	-3.31×10^{-1}	4.89×10^{-7}	1.68×10^8
(19)	$\bar{D}^0 K^{*-}$	0	0.464	-0.194	0.4537	0.753	-0.132	-0.318	-1.94×10^{-1}	3.58×10^{-6}	6.68×10^7
(20)	$\bar{D}^{*-} K^{*-}$	0	0.464	-0.194	0.4537	0.753	-0.132	-0.318	-1.94×10^{-1}	3.87×10^{-6}	6.18×10^7
(21)	$\bar{K}^0 D^-$	0.557	0.434	-0.0975	0.01601	0.0320	-0.00282	0.0137	-2.78×10^{-3}	5.39×10^{-6}	2.16×10^{11}
(22)	$\bar{K}^0 D^{*-}$	-0.555	0.434	-0.288	0.01504	0.0301	-0.00783	0.0124	-7.73×10^{-3}	3.05×10^{-7}	4.94×10^{11}
(23)	$\bar{K}^{*0} D^-$	0	0.464	-0.103	0.01593	0.0319	-0.00297	0.0136	-2.93×10^{-3}	3.14×10^{-6}	3.34×10^{11}
(24)	$\bar{K}^{*0} D^{*-}$	0	0.464	-0.103	0.01593	0.0319	-0.00297	0.0136	-2.93×10^{-3}	3.34×10^{-6}	3.14×10^{11}
(25)	ϕD_s^-	0	0.480	-0.0967	0.01599	0.0320	-0.00279	0.0137	-2.75×10^{-3}	1.64×10^{-6}	7.26×10^{11}
(26)	ϕD_s^{*-}	0	0.480	-0.0967	0.01599	0.0320	-0.00279	0.0137	-2.75×10^{-3}	4.23×10^{-6}	2.81×10^{11}
(27)	$\eta_c D_s^-$	1.474	0.708	0.260	0.001549	0.00310	0.000728	-0.00129	7.27×10^{-4}	7.27×10^{-3}	2.34×10^9
(28)	ψD_s^-	-0.827	0.708	0.664	0.00003624	0.0000725	0.0000435	0.0000233	4.35×10^{-5}	6.48×10^{-4}	7.34×10^{12}
(29)	$\eta_c D_s^{*-}$	0	0.668	0.285	0.0005167	0.00103	0.000265	-0.000425	2.65×10^{-4}	1.67×10^{-3}	7.67×10^{10}
(30)	ψD_s^{*-}	0	0.668	0.285	0.0005167	0.00103	0.000265	-0.000425	2.65×10^{-4}	1.33×10^{-2}	9.64×10^9

FIGURE CAPTIONS

Fig. 1. The unitarity triangle of the CKM matrix elements $V_{ub}^*V_{ud} + V_{cb}^*V_{cd} + V_{tb}^*V_{td} = 0$ in the (ρ, η) plane.

Fig. 2. Quark diagrams for B_c^- decaying into two mesons X^0 and X^- in the spectator approximation. a^{tree} is the color-favored tree diagram; b^{tree} is the color-suppressed tree diagram; a_q^{pen} and b_q^{pen} are the timelike penguin diagrams. Where $q, q_v = u, c$; $q', q'_v = d, s$, the subscripts "v" denote "vacuum".

Fig. 3. The timelike penguin diagrams for B_c^- decaying into two mesons X and Y . The dark dot denotes the W-loop, subscript "v" denote "vacuum". k^2 is the momentum squared carried by the virtual gluon, photon and Z^0 .




Fig. 1.







Fig. 2.




Fig. 3.